

GRZEGORZ BUKOWSKI*
BOGUSŁAW CHACHAJ*

SOIL ORIBATID MITES (ACARI, ORIBATIDA) IN THE PLOTS OF GRASS COLLECTION IN IHAR BYDGOSZCZ

Abstract

The population of soil oribatid mites in 14 plots of grass collection in IHAR in Bydgoszcz was investigated. Abundance of Oribatida population ranged 0.8–46.7 thousand indiv./m². Number of species of Oribatida ranged from 3 to 6 and the Shannon index was low. There were no influence of height and manner of spreading of grasses on oribatid population.

Keywords: Oribatida, grass, *Eupelops*, *Scheloribates*

Introduction

Institute of Plant Breeding and Acclimatization in Bydgoszcz gathers a collection of plants, both in the form of seeds and in the form of crops. Abundance and species composition of soil mites population is dependent on the microhabitat conditions. The effect on conditions in the soil have, among other things, plants overgrowth, the soil and the tillage treatments (Rajski 1961; Seniczak 1978).

* Department of Ecology, UTP University of Science and Technology, Kordeckiego 20, 85-225 Bydgoszcz, Poland, e-mails: bukowski@utp.edu.pl; chachaj@utp.edu.pl

The aim of this study was to establish whether the various species of grasses of different heights and a way of spreading, influence on the populations of soil mites.

Material and methods

The investigation was performed in the terrain of IHAR garden, in Myślęcinek part of Bydgoszcz. Grasses were cultivated in the plots of 1 x 1 m, under very intensive care (ground preparing, soil scarification, hand weeding, mowing).

Investigated material was taken from 14 plots sown with grasses of different heights and a way of spreading. The following species were investigated: 1. *Anthoxanthum odoratum* L., 2. *Nardus stricta* L., 3. *Deschampsia caespitosa* (L.) P. Beauv., 4. *Festuca airoides* Lam., 5. *Corynephorus canescens* (L.) P. Beauv., 6. *Molinia caerulea* (L.) Moench, 7. *Arrhenatherum elatius* (L.) P. Beauv. ex J. Presl et C. Presl, 8. *Alopecurus pratensis* L., 9. *Holcus lanatus* L., 10. *Phleum pratense* L., 11. *Sesleria tatrae* (Degen) Deyl, 12. *Agrostis capillaris* L., 13. *Calamagrostis pseudophragmites* (Hall.) Koel. and 14. *Phalaris arundinacea* L.

Soil samples of 50 cm³ each (3 cm deep) were taken in April and September of 2008 from each plot in 5 replicates. Samples were selected from the lower part (3 cm high) of plants and the soil. Generally 140 samples were studied. Oribatid mites were extracted in high gradient Tullgren funnels, conserved in 70% ethyl alcohol and determined to species level, including the juvenile stages. Names of oribatid species follow Weigmann (2006) and partly Subías (2004). Names of plant species follow Rutkowski (2005). The populations of oribatid species were characterized with the abundance (*A*) and constancy (*C*) indices, while the mite communities were compared with the Shannon *H* index (Odum 1971).

Results were statistically analyzed using a spreadsheet Microsoft Excel 2007 and Statistica 6.0 package. Basic descriptive statistics were carried out, and after a logarithmic transformation of data was performed using the statistical analysis of the U Mann-Whitney test and Tukey test, with a significance of $P < 0.05$ (Berthet, Gerard 1965; Łomnicki 2000).

Results and discussion

The abundance of mites in the investigated plots ranged from 4.5 to 50.9 thous. indiv./m². Oribatida ranged in the investigated plots from 19% to 92% of the all mites population (Tab. 1).

The observed abundance of mites (including oribatid mites) in the investigated plots was similar for the abundance of different types of grasslands found by other authors (Rajski 1961; Seniczak et al. 1993; Hubert et al. 2004).

In soils with strong anthropogenic pressure, fields and lawns the abundance of mites is low (Skubała et al. 2000; Bukowski, Chachaj 2008; Bukowski 2008; Zabrocka et al. 2006).

The Shannon index (*H*) of the population of oribatid mites in the investigated plots was low. In the most of investigated plots the population of mites prefers the soil layer than the lower part of plants (Tab 2). This tendency was less clear in the population of oribatid mites. Similar results obtained Seniczak et al. (2005) investigating the relatively dry meadows soils.

The analysis determined the statistical significance of differences between Oribatida group at most studied plots (Tab. 3).

In the investigated plots only 10 taxa of oribatid mites were found and the number of species ranged from 3 to 6 taxa. This low number of species of Oribatida was observed in habitats with strong anthropogenic pressure, or especially those, where were few dead organic matter (Seniczak et al. 2000a, 2000b; Domek et al. 2006; Sokołowska et al. 2006; Sokołowska, Seniczak 2005).

In the investigated plots relatively abundant were *Eupelops occultus* – species characteristic for meadows and pastures, *Scheloribates laevigatus* – species characteristic for meadows, *Scutovertex sculptus* – xerophilous species and *Tectocephus velatus* – eurytopic species (Chachaj, Seniczak 2005; Klimek et al. 1991).

Acari are very abundant in the meadows. There were no statistically significant relationships between the amount or way of spreading of particular species of grasses and the abundance of mites and species composition of oribatid mites.

Differences between populations of mites in the investigated plots could be caused by other factors, including quantity and quality of vegetable matter and the intensity of tillage treatments.

Table 1. Characteristics of grass, mite abundance (A in thousand indiv./m²), number of species of Oribatida (S), Shannon index (H), grass spread system: sl – slack, c – clump, st – stolons, grass height: l – low, a – average, h – high

		Grass species													
		1	2	3	4	5	6	7	8	9	10	11	12	13	14
Way of spreading		c	c	c	c	c	c	sl	sl	sl	sl	sl	sl	st	st
Heights		l	l	a	l	l	h	a	h	l	h	l	a	h	h
Total Acari	A	50.9	27.0	18.7	12.4	10.8	4.9	43.9	11.8	9.6	7.0	4.5	15.4	14.6	8.5
Oribatida	A	46.7	16.1	11.4	9.5	3.1	1.7	34.1	6.9	6.7	3.8	0.8	11.7	4.1	3.0
	S	5	5	5	6	5	5	6	5	6	6	3	4	6	3
	H	0.883	1.142	1.149	1.125	1.218	1.260	0.981	0.819	1.167	1.138	0.656	0.649	1.163	0.774
Eupelops occultus (C.L. Koch. 1835)	A	31.4	7.9	2.1	5.1	1.4	0.8	15.3	5.2	0.8	1.4	0.7	8.9	2.5	2.4
	C	90	90	80	80	70	70	90	60	40	70	50	90	80	90
Trichoribatens incisellus (Kramer. 1897)	A					1.1				0.1					
	C					40				10					
Liebstadia humerata Selinick. 1928	A	0.5	5.8	0.1	0.1	0.2	0.1	1.3	0.1						
	C	10	90	10	10	10	10	20	10						
Micreremus brevipes (Michael. 1888)	A												0.1		
	C												10		
Oppiella nova (Oudemans. 1902)	A	0.4	5.7	0.3	0.3	0.5	0.2	0.5	0.2	0.7	0.1	0.1	0.2	0.5	0.1
	C	10	70	20	20	30	30	10	30	30	10	30	30	30	10
Punctoribatens punctum (C.L. Koch. 1839)	A													0.6	
	C													10	
Scheloriabates laevigatus (C. L. Koch. 1835)	A	11.7			0.1			16.3	0.7	4.0	0.2	0.1		0.3	
	C	80			10			100	50	20	30	10		50	
Scutovertex sculptus Michael. 1879	A	1.6	1.3		1.1	0.1	0.3	0.1	0.2	0.1	1.9			0.1	
	C	30	30		20	10	30	10	30	10	40			10	
Suctobelba sp.	A					0.4									
	C					40									
Tectocephalus velatus Michael. 1880)	A	1.3	0.7	3.3	2.8	0.4	0.1	1.0	0.7	1.0	0.2	0.1	2.5	0.6	0.2
	C	40	40	80	80	30	10	50	40	20	30	20	90	30	10

1. *Anthoxanthum odoratum* L., 2. *Nardus stricta* L., 3. *Deschampsia caespitosa* (L.) P.Beauv., 4. *Festuca atiroides* Lam., 5. *Corynephorus canescens* (L.) P.Beauv., 6. *Molinia caerulea* (L.) Moench, 7. *Arrhenatherum elatius* (L.) P. Beauv. ex J. Presl et C. Presl, 8. *Alopecurus pratensis* L., 9. *Holcus lanatus* L., 10. *Phleum pratense* L., 11. *Sesleria tatrae* (Degen) Deyl, 12. *Agrostis capillaris* L., 13. *Calamagrostis pseudophragmites* (Hall.) Koel. and 14. *Phalaris arundinacea* L.

Table 2. Vertical distribution of mites in the investigated plots (indiv./100 cm³)

		Plots													
		1	2	3	4	5	6	7	8	9	10	11	12	13	14
Total Acari	P	136	58	28	22	16	5	53	17	16	7	8	28	17	12
	S	33	32	34	19	20	11	93	22	16	16	7	34	31	17
Oribatida	P	129	30	16	17	6	1	39	8	11	5	2	16	7	4
	S	26	23	22	15	4	4	75	15	11	8	1	22	7	6

P – lower part of plants, S – soil.

Table 3. Significant differences between the Oribatida on studied plots, x – at $P < 0.05$

		Plots													
		1	2	3	4	5	6	7	8	9	10	11	12	13	14
Plots	1		x	x	x	x	x	x	x	x	x	x	x	x	x
	2	x			x	x	x	x	x	x	x	x		x	x
	3	x				x	x	x			x	x		x	x
	4	x	x			x	x	x			x	x		x	x
	5	x	x	x	x		x	x	x	x		x	x		
	6	x	x	x	x	x		x	x	x			x		
	7	x	x	x	x	x	x		x	x	x	x	x	x	x
	8	x	x			x	x	x			x	x			x
	9	x	x			x	x	x			x	x			x
	10	x	x	x	x			x	x	x		x	x		
	11	x	x	x	x	x		x	x	x	x		x	x	x
	12	x				x	x	x				x	x		x
	13	x	x	x	x			x				x	x		
	14	x	x	x	x			x	x	x		x	x		

References

- Berthet P., Gerard G. 1965. A statistical study of microdistribution of Oribatei (Acari). Part I. The distribution pattern. *Oikos*, 16: 214–227.
- Bukowski G. 2008. Wpływ realizacji zieleni urządzonej w obszarze przemysłowym na strukturę populacji roztoczy (Acari) glebowych. *Fauna miast: ochronić różnorodność biologiczną w miastach*. P. Indykiewicz, L. Jerzak, T. Barczak (eds.). SAR „Pomorze”, Bydgoszcz, pp. 356–361.
- Bukowski G., Chachaj B. 2008. Wstępne obserwacje nad różnorodnością gatunkową mechowców (Acari, Oribatida) zbiorowisk łąkowych i ruderalnych dzielnicy Opławiec w Bydgoszczy. *Fauna miast: ochronić różnorodność biologiczną w miastach*. P. Indykiewicz, L. Jerzak, T. Barczak (eds.). SAR „Pomorze”, Bydgoszcz, pp. 309–312.

- Chachaj B., Seniczak S. 2005. The influence of sheep, cattle and horse grazing on soil mites (Acari) of lowland meadows. *Folia Biologica*, 531: 127–132.
- Domek K., Seniczak S., Kaczmarek S., Bukowski G. 2006. Preliminary studies on the effect of swine liquid manure fertilization on some soil mesofauna. In: *Advances in Polish Acarology*. G. Gabryś, S. Ignatowicz (eds.). SGGW, Warszawa, pp. 62–70.
- Hubert J., Kučerová A., Münzbergová Z. 2004. The comparison of oribatid mite (Acari: Oribatida) communities at various patches in the seminatural meadow. *Ekologia*, 23: 328–341.
- Klimek A., Seniczak S., Żelazna E., Dąbrowska B. 1991. Akarofauna (Acari) skarp osadników produktów odpadowych Janikowskich Zakładów Sodowych [The fauna of mites (Acari) of the escarpments of 'white seas' near Janikowo Soda Factory]. *Zeszyty Naukowe ATR w Bydgoszczy. Zootechnika*, 22: 151–165.
- Łomnicki A. 2000. *Wprowadzenie do statystyki dla przyrodników*. PWN, Warszawa.
- Odum E.P. 1971. *Fundamentals of ecology*. W.B. Saunders Co., Philadelphia.
- Rajski A. 1961. Studium ekologiczno-faunistyczne nad mechowcami (Acari, Oribatei) w kilku zespołach roślinnych. I. *Ekologia Pr. Kom. Biol. PZPN*, 25: 1–160.
- Rutkowski L. 2005. *Klucz do oznaczania roślin naczyniowych Polski niżowej*. Wydawnictwo Naukowe PWN, Warszawa.
- Seniczak S. 1978. *Stadia młodociane mechowców (Acari, Oribatei) jako istotny składnik zgrupowań tych roztoczy przetwarzających glebową substancję organiczną*. Uniwersytet Mikołaja Kopernika, Toruń.
- Seniczak S., Bukowski G., Kaczmarek S., Bukowska H., Kobierski M. 2000a. The influence of different plant cover of forest dune on soil mites (Acari) in the Tuchola National Park. In: *Akarologia polska u progu nowego tysiąclecia*. S. Ignatowicz (ed.). SGGW, Warszawa, pp. 202–210.
- Seniczak S., Chachaj B., Wasieńska B., Graczyk R., Kwiatkowski J., Waldon B., Kobierski M. 2005. Wpływ nawożenia łąki wodą amoniakalną z oczyszczalni ścieków na roztocze glebowe (Acari). *Zeszyty Naukowe ATR w Bydgoszczy, Zootechnika*, 35: 17–26.
- Seniczak S., Dąbrowski J., Dąbrowska B., Gackowski G. 1993. Wpływ aerozoli ścieków pochodzących z Zakładów Chemicznych „Organika-Zachem” w Bydgoszczy na akarofaunę łąkową, ze szczególnym uwzględnieniem Oribatida (Acari). *Zeszyty Naukowe ATR w Bydgoszczy, Zootechnika*, 23: 121–132.
- Seniczak S., Kaczmarek S., Zaręba J., Seniczak A., Marcysiak K., Kobierski M. 2000b. The preliminary observations on soil mites (Acari) of lawns in the neighbourhood of busy road in Bydgoszcz. In: *Akarologia polska u progu nowego tysiąclecia*. S. Ignatowicz (ed.). SGGW, Warszawa, pp. 211–219.

- Skubała P., Madej G., Krajczyk A. 2000. Mites (Acari: Oribatida, Mesostigmata) of agrocenoses subjected to pollution. In: *Akarologia polska u progu nowego tysiąclecia*. S. Ignatowicz (ed.). SGGW, Warszawa, pp. 136–142.
- Sokołowska L., Seniczak S. 2005. Wpływ nawożenia gnojówką bydlęcą na wydajność przemianowego użytku zielonego i liczebność wybranych grup mezofauny glebowej. *Zeszyty Naukowe ATR Bydgoszcz, Zootechnika*, 35: 55–67.
- Sokołowska L., Seniczak S., Kaczmarek S., Bukowski G. 2006. Preliminary studies on the effect of cattle slurry fertilization on some soil mesofauna groups. In: *Advances in Polish Acarology*. G. Gabryś, S. Ignatowicz (ed.). SGGW, Warszawa, pp. 399–409.
- Subías L.S. 2004. Listado sistemático, sinonímico y biogeográfico de los Ácaros Oribátidos (Acariformes, Oribatida) del mundo (1748–2002). *Graellsia*, 60: 3–305.
- Weigmann G. 2006. *Hornmilben (Oribatida)*. Die Tierwelt Deutschlands. Vol. 76.
- Zabrocka E., Seniczak S., Seniczak A., Bukowski G., Dudek S. 2006. Wpływ nawadniania i nawożenia mineralnego na jakość plonu trawy i roztocze (Acari) glebowe. *Zeszyty Naukowe ATR Bydgoszcz, Zootechnika*, 36: 51–58.

MECHOWCE (ACARI, ORIBATIDA) GLEBOWE NA POLETKACH KOLEKCJI TRAW IHAR W BYDGOSZCZY

Streszczenie

Zbadano populacje mechowców glebowych na 14 poletkach kolekcji traw IHAR w Bydgoszczy. Liczebność populacji Oribatida wahała się od 0,8 do 46,7 tysięcy osobn./m². Liczba gatunków Oribatida wahała się od 3 do 6, a indeks Shannona był niski. Nie stwierdzono wpływu wysokości i sposobu krzewienia się traw na populacje Oribatida.

Słowa kluczowe: mechowce, trawy, *Eupelops*, *Scheloribates*

Cite this article as: Bukowski G., Chachaj B. 2015. Soil oribatid mites (Acari, Oribatida) in the plots of grass collection in IHAR Bydgoszcz. *Acta biologica*, 22: 63–69.

